$pp \to J/\psi + \Upsilon + X$ as a clean probe to the quarkonium production mechanism

P. Ko* and Chaehyun Yu†
School of Physics, KIAS, Seoul 130-722, Korea

Jungil Lee‡
Department of Physics, Korea University, Seoul 136-701, Korea

We report that, unlike most inclusive quarkonium-production processes, the production rate for $pp \to J/\psi + \Upsilon + X$ dominantly depends on two not well-known long-distance nonrelativistic QCD matrix elements, $\langle O_8^{J/\psi}(^3S_1)\rangle$ and $\langle O_8^{\Upsilon}(^3S_1)\rangle$ at leading order in the strong coupling constant, which account for the transition probabilities of the color-octet spin-triplet heavy-quark-antiquark pairs $c\bar{c}_8(^3S_1) \to J/\psi$ and $b\bar{b}_8(^3S_1) \to \Upsilon$, respectively. With the integrated luminosity $\sim 100\,\mathrm{fb}^{-1}$ at the center-of-momentum energy $\sqrt{s} = 14\,\mathrm{TeV}$ we expect that approximately 1900 $pp \to J/\psi + \Upsilon + X$ events can be observed at the CERN Large Hadron Collider by tagging muon pairs, which are enough to probe to the color-octet mechanism. The forthcoming measurement may provide a useful constraint to resolve the decade-old puzzle for the polarization of prompt J/ψ at the Fermilab Tevatron. If corresponding measured rate is significantly less than the prediction, it may imply that the current values for the color-octet matrix elements are overestimated.

PACS numbers: 12.38.-t, 13.85.Ni, 14.40.Pq

arXiv:1006.3846v2 [hep-ph] 14 Mar 2011

As an effective field theory of QCD, the nonrelativistic QCD (NRQCD) factorization approach [1] has achieved great progress in understanding the production and decay mechanism of heavy quarkonia [2]. The factorization has been proved for the electromagnetic and light-hadronic decays [1] and also for a few specific exclusive production processes very recently [3–5]. The factorization conjecture for the inclusive production has been employed to resolve the large surplus of $\psi(2S)$ and prompt J/ψ at the Fermilab Tevatron [6], which lead to a remarkable prediction that the prompt J/ψ must be transversely polarized at the large transverse momentum (p_T) [7]. However, the empirical data for the polarization of prompt J/ψ and $\psi(2S)$ measured by the CDF Collaboration [8, 9] are in disagreement with these predictions. That is what we call the puzzle for the polarization of prompt J/ψ at the Tevatron.

These predictions strongly depend on the determination of the long-distance NRQCD matrix element $\langle O_n^H(^{2s+1}L_J)\rangle$, which accounts for the probability of the heavy-quark-antiquark pair $Q\bar{Q}$ with the spectroscopic state $^{2s+1}L_J$ to evolve into a heavy quarkonium H. The color-singlet NRQCD matrix elements $\langle O_1^H(^3S_1)\rangle$ for $H = J/\psi$ and Υ are well known through precisely measured values for their leptonic decay rates. On the other hand, the color-octet counterparts such as $\langle O_8^H(^3S_1)\rangle$, $\langle O_8^H(^1S_0)\rangle$, and $\langle O_8^H(^3P_J)\rangle$ for $J = 0$, 1, 2 are not determined accurately, because in most of the hadroproduction rates for H all of these color-octet matrix elements involve simultaneously and the p_T spectra are not well distinguishable to be fit to the data [10, 11]. To make the situation worse, large next-to-leading-order corrections dramatically modify the p_T spectra of these processes [12–15]. As a result, the matrix elements determined from various processes are not consistent with each other [15–21]. In particular, the actual values for the color-octet matrix elements $\langle O_8^{J/\psi}(^1S_0)\rangle$ and $\langle O_8^{J/\psi}(^3P_0)\rangle$ might be significantly smaller than the naive estimates based on the velocity-scaling rules of NRQCD [21]. In this situation, it would be highly desirable to find a hadroproduction process that involves fewer numbers of color-octet channels.

In this paper, we propose that $pp \to J/\psi + \Upsilon + X$ at the CERN Large Hadron Collider (LHC) could be a clean probe to the color-octet mechanism of NRQCD and compute its p_T spectrum and the total production rate at the center-of-momentum energy $\sqrt{s} = 14\,\mathrm{TeV}$. Unlike most inclusive quarkonium-production processes [7, 22–24], the production rate for $pp \to J/\psi + \Upsilon + X$ dominantly depends on two not well-known long-distance factors $\langle O_8^{J/\psi}(^3S_1)\rangle$ and $\langle O_8^{\Upsilon}(^3S_1)\rangle$ and the color-singlet contribution is suppressed at leading order (LO) in the strong coupling constant α_s in the asymptotic limit $p_T \to \infty$. The forthcoming measurement may provide a useful constraint to resolve the decade-old puzzle for the polarization of prompt J/ψ at the Tevatron.

This paper is organized as follows. We first summarize the NRQCD factorization formula for $pp \to J/\psi + \Upsilon + X$ and describe a systematic way to determine the dominant contributions. Next we provide detailed arguments to convince the dominance of $\langle O_8^{J/\psi}(^3S_1)\rangle$ and $\langle O_8^{\Upsilon}(^3S_1)\rangle$ contributions in the p_T spectrum at LO in α_s and the typical velocity v_Q of the heavy quark Q in the quarkonium rest frame. Our prediction for the production rate and p_T spectrum at the LHC follows and then we conclude.

The NRQCD factorization formula for the differential cross section $d\sigma$ of $pp \to J/\psi + \Upsilon + X$ has the following schematic form:

$$d\sigma = f_{i/p} \otimes f_{j/p} \otimes d\hat{\sigma}_{ij \to c\bar{c}_{n_1} + b\bar{b}_{n_2}} \langle O_{n_1}^{J/\psi}\rangle \langle O_{n_2}^{\Upsilon}\rangle, \quad (1)$$




FIG. 1. Typical Feynman diagrams for gluon-initiated parton processes for $pp \to J/\psi + \Upsilon + X$ at order α_s^4.




FIG. 2. Typical Feynman diagrams for the color-singlet contribution to $pp \to J/\psi + \Upsilon + X$ at order α_s^6, which is leading.

where $f_{i/p}$ is the parton distribution function (PDF), $d\hat{\sigma}$ is the parton-level cross section, which is perturbatively calculable in powers of α_s, $\langle O_{n_i}^{H_i} \rangle$ is the NRQCD matrix element for the quarkonium H_i involving the NRQCD four-quark operator $O_{n_i}^{H_i}$ [1]. The subscript n_i represents the spectroscopic state of the $Q\bar{Q}$ pair. For example, $Q\bar{Q}_{n_i} = Q\bar{Q}_n(^{2s+1}L_J)$ stands for the color-singlet ($n = 1$) or -octet ($n = 8$) $Q\bar{Q}$ pair with the spin s, the orbital angular momentum L, and the total angular momentum J for $Q = c$ or b. The symbol $\otimes$ indicates the convolution over the partons' longitudinal momentum fractions and the summation over partons i and j is assumed. According to the velocity-scaling rules of NRQCD [1], the production rates for the spin-triplet S-wave quarkonium through the color-octet states $Q\bar{Q}_8(^3S_1)$, $Q\bar{Q}_8(^1S_0)$, and $Q\bar{Q}_8(^3P_J)$ for $J = 0$, 1, 2 are suppressed by v_Q^4, v_Q^3, and v_Q^4, respectively, compared to the color-singlet state $Q\bar{Q}_1(^3S_1)$. Let us call the product of these factors for a process the "velocity-scaling factor" $\mathcal{V}$. Therefore, Eq. (1) is a power series in α_s, v_c, and v_b.

At LO in α_s, only gg fusion and $q\bar{q}$ annihilation contribute to the parton processes $ij \to c\bar{c}_{n_1} + b\bar{b}_{n_2}$ of order α_s^4 and the channels with $ij(ji) = gq$ and $g\bar{q}$ are missing. At the center-of-momentum energy of the LHC one probes the small-x region of the PDF, where the gluon contribution dominates over the quark contents. Therefore, we consider only the gg initial states. Typical Feynman diagrams for the gluon-initiated parton process $gg \to c\bar{c}_{n_1} + b\bar{b}_{n_2}$ are shown in Fig. 1, where at least one $Q\bar{Q}$ pair is in a color-octet state. Leading color-singlet contribution $c\bar{c}_1(^3S_1) + b\bar{b}_1(^3S_1)$ appears only at order α_s^6, which are shown in Fig. 2.

We first consider the color-octet contribution $c\bar{c}_8(^3S_1) + b\bar{b}_8(^3S_1)$ in Figs. 1 (a)–(d) with $\mathcal{V} = v_c^4 v_b^4$, to which 36 Feynman diagrams contribute. At large p_T the double-fragmentation contribution [Fig. 1 (a)] dominates because of the kinematic enhancement. Figure 1 (b) also represents mixed contributions $c\bar{c}_8(^3S_1) + b\bar{b}_1(^3S_1)$ with $\mathcal{V} = v_c^4$ and $c\bar{c}_1(^3S_1) + b\bar{b}_8(^3S_1)$ with $\mathcal{V} = v_b^4$, each of which has 6 Feynman diagrams. If p_T is not large enough, then these mixed contributions must dominate over $c\bar{c}_8(^3S_1) + b\bar{b}_8(^3S_1)$ by the enhancement factors $1/v_b^4$ or $1/v_c^4$ while the double-fragmentation contribution dominates at large p_T.

In addition, $c\bar{c}_8(^3S_1) + b\bar{b}_8(^{2s+1}L_J)$ and $c\bar{c}_8(^{2s+1}L_J) + b\bar{b}_8(^3S_1)$, where $^{2s+1}L_J = {}^1S_0$ or ^3P_J, contribute to diagrams in Figs. 1 (b)–1 (d), with $\mathcal{V} = v_c^\alpha v_b^\beta$. Here, α, $\beta = 3$ for $^{2s+1}L_J = {}^1S_0$ and 4 for ^3P_J. The $\mathcal{V}$ is comparable to that of the double-fragmentation, while they are suppressed at least by either v_c^3 or v_b^3 in comparison with $c\bar{c}_1(^3S_1) + b\bar{b}_1(^3S_1)$. Although the single-fragmentation channel in Fig. 1 (c) may grow up at large p_T, that contribution is dominated by the double-fragmentation [Fig. 1 (a)] by a factor of $(m_c/p_T)^4$ or $(m_b/p_T)^4$. Hence, it is consistent to ignore $c\bar{c}_8(^3S_1) + b\bar{b}_8(^{2s+1}L_J)$ and $c\bar{c}_8(^{2s+1}L_J) + b\bar{b}_8(^3S_1)$ with $^{2s+1}L_J = {}^1S_0$ or ^3P_J over the whole p_T range.

As the last color-octet contribution, there are $c\bar{c}_8(^{2s+1}L_J) + b\bar{b}_8(^{2s'+1}L'_{J'})$, where $^{2s+1}L_J$, $^{2s'+1}L'_{J'} = {}^1S_0$ or ^3P_J, whose typical Feynman diagram is shown in Fig. 1 (d). In this case $\mathcal{V} = v_c^\alpha v_b^\beta$ with α, $\beta = 3$ for $^{2s+1}L_J = {}^1S_0$ and 4 for ^3P_J. Thus the contributions are suppressed compared to the color-singlet one. Because they do not have any fragmentation contributions, they are dominated by the double-fragmentation [Fig. 1 (a)] by $(m_c/p_T)^4(m_b/p_T)^4$ in the large-p_T region.

So far, we have shown that, at LO in α_s and v_Q, it is sufficient to consider only $c\bar{c}_8(^3S_1) + b\bar{b}_8(^3S_1)$, $c\bar{c}_8(^3S_1) + b\bar{b}_1(^3S_1)$, and $c\bar{c}_1(^3S_1) + b\bar{b}_8(^3S_1)$ to describe the p_T spectrum of $pp \to J/\psi + \Upsilon + X$. The parton-level differential cross sections for these processes are available in Ref. [25].

Based on the formalism that we have described earlier, we are ready to carry out the numerical calculation of the production rates for $J/\psi + \Upsilon + X$ at the LHC. The formula (1) involves the factorization of the PDF and the parton-level cross section $d\hat{\sigma}$ with the factorization scale μ. For the scale μ, we take the transverse mass $\mu = m_T = (4m_Q^2 + p_T^2)^{1/2}$ with the heavy-quark masses $m_c = 1.5$ GeV and $m_b = 4.7$ GeV. At LO in α_s on which we are working, there are no additional hard jets and, therefore, p_T and m_T are defined unambiguously. We employ the CTEQ6L parametrization [26] for the PDF. We evaluate α_s by setting the renormalization scale to be m_T and use the next-to-leading-order formula for the running coupling constant to be consistent with Ref. [26].

In order to evaluate the production rate (1), we have to know the values for the NRQCD matrix elements $\langle O_n^H(^3S_1) \rangle$ for $H = J/\psi$ and Υ, where $n = 1$ or 8. The color-singlet matrix element $\langle O_1^H(^3S_1) \rangle$ is usually determined from the leptonic decay rate of H, which is the most precisely measured value involving H. We quote $\langle O_1^{J/\psi}(^3S_1) \rangle = 1.32\,\mathrm{GeV}^3$ [27] and $\langle O_1^{\Upsilon}(^3S_1) \rangle = 9.21\,\mathrm{GeV}^3$ [28]. The color-octet matrix element $\langle O_8^{J/\psi}(^3S_1) \rangle$ has been fit to the p_T spectrum of the




FIG. 3. The differential cross section $d\sigma/dp_T$ for $pp \to J/\psi + \Upsilon + X$ at $\sqrt{s}$ =14 TeV in units of pb/GeV as a function of p_T integrated over the rapidity range $|y| < 2.4$. The solid, dashed, dashed-dotted, and dotted curves represent the total, $c\bar{c}_1(^3S_1) + b\bar{b}_8(^3S_1)$, $c\bar{c}_8(^3S_1) + b\bar{b}_1(^3S_1)$, and $c\bar{c}_8(^3S_1) + b\bar{b}_8(^3S_1)$ contributions, respectively.

inclusive prompt J/ψ production rate at the Tevatron in the large-p_T region. The matrix element $\langle O_8^{\Upsilon}(^3S_1)\rangle$ has been fit [11] to the Tevatron data and used for the polarization analysis [29]. Various determinations of these matrix elements can be found, for example, in Ref. [30]. In this work, we use $\langle O_8^{J/\psi}(^3S_1)\rangle = 3.9 \times 10^{-3}$ GeV^3 [7], $\langle O_8^{\Upsilon}(^3S_1)\rangle = 1.5 \times 10^{-1}$ GeV^3 [30].

The prediction for $d\sigma/dp_T$ of $pp \to J/\psi + \Upsilon + X$ at $\sqrt{s}$ =14 TeV integrated over the rapidity range $|y| < 2.4$ is shown as the solid curve in Fig. 3. $d\sigma/dp_T$ vanishes at $p_T = 0$ and increases rapidly until it reaches the maximum value $d\sigma/dp_T|_{\rm max} = 2.9$ pb/GeV at $p_T = 1.5$ GeV. Then it monotonically decays as p_T increases. Near $p_T = 0$, $c\bar{c}_1(^3S_1)+b\bar{b}_8(^3S_1)$ dominates and it is the largest for $p_T \lesssim 4$ GeV. At large p_T, the double-fragmentation contribution $c\bar{c}_8(^3S_1) + b\bar{b}_8(^3S_1)$ [Fig. 1 (a)] dominates and it is the largest for $p_T \gtrsim 6$ GeV. In the remaining region, 4 GeV $\lesssim p_T \lesssim$ 6 GeV, all of the three contributions compete together.

As is mentioned earlier, we have ignored the color-singlet contribution in Fig. 2 because it is suppressed compared to the color-octet one by a relative order of α_s^2. In order to assure that this approximation is safe, we make a rough estimate of the color-singlet contribution to $pp \to J/\psi + \Upsilon + X$. The corresponding parton processes consist of 3 types: Types A, B, and C are shown in Fig. 2 (a)–2 (c), respectively.

We first classify the scaling behavior of the amplitude for the double-fragmentation diagram shown in Fig. 1 (a). The propagator for the exchanged gluon has the scaling $1/p_T^2$ and the gluon propagators attached to the $c\bar{c}$ and $b\bar{b}$ pairs are of order $1/m_c^2$ and $1/m_b^2$, respectively. The product of the two triple-gluon vertices must scale as the typical momentum squared p_T^2. Therefore, the resultant relative scaling of the double-fragmentation process is $v_c^4 v_b^4/(m_c^4 m_b^4)$, where we have included the velocity-scaling factor $\mathcal{V} = v_c^4 v_b^4$.

In the case of type A, there are four heavy-quark propagators and one gluon propagator which have a large momentum transfer that accounts for the scaling factor $1/p_T^6$. The phase space for the $c\bar{c}_1 + b\bar{b}_1 + gg$ final state enhances the scaling factor by an order of p_T^4 in comparison with that for the two-body final state $c\bar{c}_8 + b\bar{b}_8$ of the double-fragmentation process. As a result, the type A process is scaled by $1/p_T^8$ compared to the double-fragmentation process. In a similar manner, the scaling factor for type B is the same as that of type A. In types A and B, there are extra hard jets in the final states. Therefore, the color-octet contribution, which does not have hard jets, is distinguished from these color-singlet contributions that can simply be removed by imposing an appropriate veto. Type C diagrams involve finite box diagrams with two virtual gluons whose momentum must be of order the typical momentum transfer p_T. A rough estimate of the scaling can be found by substituting a typical momentum transfer p_T to the two gluon propagators, four heavy-quark propagators, and the measure of the loop momentum. The relative scaling of type C process is again $1/p_T^8$.

Therefore, the suppression factor for the color-singlet contribution relative to the color-octet double-fragmentation process is $[\alpha_s/(4\pi)]^2[m_c/(v_c p_T)]^4 \times [m_b/(v_b p_T)]^4$ where we have included the strong coupling suppression factor α_s^2 to the color-singlet contribution. However, if p_T is small, then the factors p_T in this scaling should be of order m_Q and, therefore, the suppression factor becomes roughly $\alpha_s^2/[(4\pi)^2 v_b^4]$ in comparison with $c\bar{c}_1(^3S_1) + b\bar{b}_8(^3S_1)$, which dominates over the double-fragmentation contribution as $p_T \to 0$. This factor is of order 10^{-2} if we assume $v_b^2 \sim 0.1$ and we use the renormalization scale of order m_b. According to this rough estimate, we expect that our prediction may not be contaminated by an order-α_s^6 color-singlet contribution even at small p_T. If we introduce a lower p_T cut $p_T \gtrsim 5$ GeV, where the double-fragmentation rises up, then the color-octet contribution becomes more significant. In addition, by imposing a veto that the final state must not include hard jets, one can remove the color-singlet contribution of types A and B, which are potential sources of a large background.

The kinematic enhancement of the double gluon fragmentation can also be applied to the color-singlet processes of order α_s^8 through the subprocesses $gg, q\bar{q} \to gg$ followed by the fragmentations of the two final-state gluons into $c\bar{c}_1(^3S_1) + gg$ and $b\bar{b}_1(^3S_1) + gg$, respectively. This process is suppressed compared to the color-octet double gluon fragmentation $g \to c\bar{c}_8(^3S_1)$ and $g \to b\bar{b}_8(^3S_1)$ that we have calculated in this paper by a factor

of order $\alpha_s^4/[(4\pi)^4 v_c^4 v_b^4] \sim 7\times 10^{-5}$ with an assumption $\alpha_s = 0.2$ at threshold with the scale $\mu = 2m_Q$.[1] In addition, one can consider the mixed double gluon fragmentation of order α_s^6 through the subprocesses $gg,\, q\bar{q} \to gg$ followed by the fragmentations of the two final-state gluons into $c\bar{c}_8(b\bar{b}_8)(^3S_1)$ and $b\bar{b}_1(c\bar{c}_1)(^3S_1) + gg$, respectively. These mixed double-fragmentation contributions are also suppressed relative to the color-octet counterpart by factors of $\alpha_s^2/[(4\pi)^2 v_b^4] \sim 3\times 10^{-2}$ and $\alpha_s^2/[(4\pi)^2 v_c^4] \sim 3\times 10^{-3}$ for the $c\bar{c}_8 + b\bar{b}_1$ and $b\bar{b}_8 + c\bar{c}_1$ fragmentations, respectively.[2] In summary, the mixed double-fragmentation contributions may occupy only a few percents of the color-octet double fragmentation and the pure color-singlet double-fragmentation contribution seems to be well suppressed compared to the color-octet double-fragmentation contribution.

Based on the NRQCD factorization formalism, we have computed the p_T spectrum $d\sigma/dp_T$ and the total cross section $\sigma_{\rm tot}$ for $pp \to J/\psi + \Upsilon + X$. The short-distance coefficients were computed at LO in α_s and v_Q. $\sigma_{\rm tot}$ integrated over the rapidity and the transverse-momentum range $|y| < 2.4$ and $p_T < 30\,{\rm GeV}$ is predicted to be $\sigma_{\rm tot} = 13\,{\rm pb}$ at $\sqrt{s} = 14\,{\rm TeV}$. Because the production rate is subject to the numerical values for $\langle O_8^{J/\psi}(^3S_1)\rangle$ and $\langle O_8^{\Upsilon}(^3S_1)\rangle$, the forthcoming empirical rate may provide information whether the values were reasonable or not.

The feature of this LO process that the color-singlet channel is suppressed provides us with a clean probe to the color-octet mechanism. In comparison with the inclusive single-quarkonium production at the hadron colliders, the suppression of the color-singlet channel is more significant by a relative factor of $[\alpha_s/(4\pi)]^2[m_Q/(v_Q p_T)]^4$ at large p_T. The p_T spectrum is dominated by $c\bar{c}_1(^3S_1) + b\bar{b}_8(^3S_1)$ and $c\bar{c}_8(^3S_1) + b\bar{b}_8(^3S_1)$ channels at the low- and large-p_T region, respectively, and the $c\bar{c}_8(^3S_1) + b\bar{b}_1(^3S_1)$ contribution is comparable to those two listed above only around the region $4\,{\rm GeV} \lesssim p_T \lesssim 6\,{\rm GeV}$. Assuming the integrated luminosity $\sim 100\,{\rm fb}^{-1}$ at $\sqrt{s}$ =14 TeV and considering the branching fractions $B[J/\psi \to \mu^+\mu^-] = 5.93\,\%$ and $B[\Upsilon \to \mu^+\mu^-] = 2.48\,\%$ [32], we expect that approximately 1900, 520, and 160 events can be observed by tagging muon pairs under the cuts $p_T \geq 0$, 5, and 10 GeV, respectively, at the LHC. If one can improve the acceptances for J/ψ and Υ by extensive Monte Carlo studies of final-muon pairs, then the observation of the events can be quite promising in the near future. Inclusion of e^+e^- decay modes of J/ψ and Υ may increase the number of events by a factor of 4. One can also improve the prediction by including the subprocess $q\bar{q} \to J/\psi + \Upsilon$ via two-gluon exchange that is neglected in this work. If the forthcoming measured rate is significantly less than our prediction, then it may be an indication that the current values for the octet matrix elements are overestimated.

We consider possible sources that may contaminate our predictions. In this work, we have computed the rate for the direct J/ψ's rather than prompt J/ψ's that include feeddowns from higher resonances and that exclude those from the B-meson decay. While nonprompt signals from B decays can be eliminated with the aid of silicon vertex detectors, the separation of the feeddowns from higher resonances is not easy. Therefore, the simultaneous determination of the color-octet NRQCD matrix elements $\langle O_8^{J/\psi}(^3S_1)\rangle$ and $\langle O_8^{\Upsilon}(^3S_1)\rangle$ using this process may be nontrivial. However, the dominance of the $c\bar{c}_8(^3S_1) + b\bar{b}_8(^3S_1)$ contribution may still be true, even after including the feeddowns from higher resonances, especially at large p_T due to the double-fragmentation process. Once the p_T distribution for the $J/\psi + \Upsilon$ production is measured, it may provide us with significant constraints on the product of the two matrix elements.

We note that the total rate and the p_T spectrum for the inclusive single-quarkonium production at hadron colliders suffer from large next-to-leading-order corrections in α_s. In particular, the next-to-leading-order corrections to the color-singlet contribution to the inclusive J/ψ production enhance the rate by an order of magnitude [12–14]. Therefore, it is natural to worry that the inclusion of order-α_s^6 contributions to this process may modify the p_T spectrum significantly. However, in the case of $J/\psi + \Upsilon$ production, the suppression factor of the color-singlet channel to the color-octet one is α_s^2/p_T^8 which is significantly smaller than the corresponding factor $1/p_T^4$ for the single-quarkonium production, in which the color-singlet and color-octet contributions are of the same order (α_s^3) at LO. We anticipate that our arguments can be tested quantitatively by measurements and by explicit calculations of complete order-α_s^6 contributions to the color-singlet channel in near future. Once the suppression of the color-singlet channel is confirmed at that order, then the measurement of the p_T spectrum of $J/\psi + \Upsilon$ at the LHC may play an important role to determine the color-octet NRQCD matrix elements $\langle O_8^{J/\psi}(^3S_1)\rangle$ and $\langle O_8^{\Upsilon}(^3S_1)\rangle$. Even in the worst case, that one could not observe the p_T spectrum at the

[1] Our order-of-magnitude estimate relies somewhat on the extra factors of (4π), which are from the phase space factor or loop integration. It is well known that the inclusion of the factor (4π) improves the naive estimates, for example, in the muon decay rate using the Fermi's four-fermion interaction Lagrangian for charged weak interaction. Our naive estimates may break down if there are large enhancement factors in the color-singlet channels. An explicit calculation of the complete color-singlet contribution may reveal if our rough power counting is indeed reasonable.

[2] If we make use of the fragmentation probability at threshold given in Ref. [31], we can make rough estimates on the ratios of the double-fragmentation processes $c\bar{c}_1 + b\bar{b}_1$, $c\bar{c}_8 + b\bar{b}_1$, and $b\bar{b}_8 + c\bar{c}_1$ compared to the $c\bar{c}_8 + b\bar{b}_8$ fragmentation as $\sim 2\times 10^{-4}$, $\sim 5\times 10^{-3}$, and $\sim 4\times 10^{-2}$, respectively. These values are consistent with the scaling factors listed above.

desired level, at least one can obtain strong upper bounds to both NRQCD matrix elements, that were not easy to extract empirically in other experiments. Therefore, the forthcoming measurement may provide useful information that can be used to understand the mechanism of the decade-old puzzle for the polarization of prompt J/ψ at the Tevatron.

In summary, we have computed the production rate and p_T distribution for $pp \to J/\psi + \Upsilon + X$ at the LHC in the framework of the NRQCD factorization formalism at leading order in α_s. Unlike most inclusive single-quarkonium-production processes, in which at least three independent color-octet NRQCD matrix elements $\langle O_8^H(^3S_1)\rangle$, $\langle O_8^H(^1S_0)\rangle$, and $\langle O_8^H(^3P_J)\rangle$ involve, this process depends dominantly on two matrix elements $\langle O_8^{J/\psi}(^3S_1)\rangle$ and $\langle O_8^{\Upsilon}(^3S_1)\rangle$ at LO in α_s, especially at large p_T. If the corresponding measured rate is significantly less than the prediction presented in this work, it may imply that current values for the octet matrix elements are overestimated. The forthcoming measurement of p_T distribution may probe to the color-octet mechanism in quarkonium production, by imposing strong constraints to these NRQCD matrix elements.

The authors would like thank Kuang-Ta Chao and Rong Li for their useful comments. They also thank Suyong Choi for providing valuable information regarding four-muon analysis at the CMS experiment. A part of the work (P.K.) was done at Aspen Center for Physics during the summer workshop in 2010. The work of P.K. was supported in part by the National Research Foundation (NRF) through the Korea Neutrino Research Center (KNRC) at Seoul National University. This work was supported by the Basic Science Research Program through the NRF of Korea funded by the MEST under Contracts No. 2009-0072689 (C.Y.) and No. 2010-0000144 (J.L.).

* pko@kias.re.kr
† chyu@kias.re.kr
‡ jungil@korea.ac.kr

[1] G. T. Bodwin, E. Braaten, and G. P. Lepage, Phys. Rev. D **51**, 1125 (1995) [Erratum-ibid. D **55**, 5853 (1997)] [arXiv:hep-ph/9407339].
[2] See, for example, N. Brambilla *et al.* (Quarkonium Working Group), CERN Report No. CERN-2005-005 (2005) [arXiv:hep-ph/0412158].
[3] G. T. Bodwin, X. Garcia i Tormo, and J. Lee, Phys. Rev. Lett. **101**, 102002 (2008) [arXiv:0805.3876 [hep-ph]].
[4] G. T. Bodwin, X. Garcia i Tormo, and J. Lee, Phys. Rev. D **81**, 114005 (2010) [arXiv:0903.0569 [hep-ph]].
[5] G. T. Bodwin, X. Garcia i Tormo, and J. Lee, Phys. Rev. D **81**, 114014 (2010) [arXiv:1003.0061 [hep-ph]].
[6] E. Braaten and S. Fleming, Phys. Rev. Lett. **74**, 3327 (1995) [arXiv:hep-ph/9411365].
[7] E. Braaten, B. A. Kniehl, and J. Lee, Phys. Rev. D **62**, 094005 (2000) [arXiv:hep-ph/9911436].
[8] A. A. Affolder *et al.* (CDF Collaboration), Phys. Rev. Lett. **85**, 2886 (2000) [arXiv:hep-ex/0004027].
[9] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. Lett. **99**, 132001 (2007) [arXiv:0704.0638 [hep-ex]].
[10] P. Cho and A. K. Leibovich, Phys. Rev. D **53**, 6203 (1996) [arXiv:hep-ph/9511315].
[11] E. Braaten, S. Fleming, and A. K. Leibovich, Phys. Rev. D **63**, 094006 (2001) [arXiv:hep-ph/0008091].
[12] J. M. Campbell, F. Maltoni, and F. Tramontano, Phys. Rev. Lett. **98**, 252002 (2007) [arXiv:hep-ph/0703113].
[13] B. Gong and J. X. Wang, Phys. Rev. Lett. **100**, 232001 (2008) [arXiv:0802.3727 [hep-ph]]; Phys. Rev. D **78**, 074011 (2008) [arXiv:0805.2469 [hep-ph]].
[14] P. Artoisenet, J. M. Campbell, J. P. Lansberg, F. Maltoni, and F. Tramontano, Phys. Rev. Lett. **101**, 152001 (2008) [arXiv:0806.3282 [hep-ph]].
[15] B. Gong, X. Q. Li, and J. X. Wang, Phys. Lett. B **673**, 197 (2009) [arXiv:0805.4751 [hep-ph]].
[16] P. Ko, J. Lee, and H. S. Song, Phys. Rev. D **54**, 4312 (1996) [Erratum-ibid. D **60**, 119902 (1999)] [arXiv:hep-ph/9602223].
[17] M. Klasen, B. A. Kniehl, L. N. Mihaila, and M. Steinhauser, Phys. Rev. Lett. **89**, 032001 (2002) [arXiv:hep-ph/0112259].
[18] P. Artoisenet, J. M. Campbell, F. Maltoni, and F. Tramontano, Phys. Rev. Lett. **102**, 142001 (2009) [arXiv:0901.4352 [hep-ph]];
[19] C. H. Chang, R. Li, and J. X. Wang, Phys. Rev. D **80**, 034020 (2009) [arXiv:0901.4749 [hep-ph]].
[20] M. Butenschoen and B. A. Kniehl, Phys. Rev. Lett. **104**, 072001 (2010) [arXiv:0909.2798 [hep-ph]].
[21] Y. J. Zhang, Y. Q. Ma, K. Wang, and K. T. Chao, Phys. Rev. D **81**, 034015 (2010) [arXiv:0911.2166 [hep-ph]].
[22] C. F. Qiao, Phys. Rev. D **66**, 057504 (2002) [arXiv:hep-ph/0206093].
[23] R. Li, Y. J. Zhang, and K. T. Chao, Phys. Rev. D **80**, 014020 (2009) [arXiv:0903.2250 [hep-ph]].
[24] C. F. Qiao, L. P. Sun, and P. Sun, J. Phys. G **37**, 075019 (2010) [arXiv:0903.0954 [hep-ph]].
[25] P. Ko, C. Yu, and J. Lee, JHEP **1101**, 070 (2011) [arXiv:1007.3095 [hep-ph]].
[26] J. Pumplin, D. R. Stump, J. Huston, H. L. Lai, P. M. Nadolsky, and W. K. Tung, JHEP **0207**, 012 (2002) [arXiv:hep-ph/0201195].
[27] G. T. Bodwin, D. Kang, and J. Lee, Phys. Rev. D **74**, 014014 (2006) [arXiv:hep-ph/0603186]; G. T. Bodwin, H. S. Chung, D. Kang, J. Lee, and C. Yu, Phys. Rev. D **77**, 094017 (2008) [arXiv:0710.0994 [hep-ph]].
[28] D. Kang, T. Kim, J. Lee, and C. Yu, Phys. Rev. D **76**, 114018 (2007) [arXiv:0707.4056 [hep-ph]]; H. S. Chung, T. Kim, and J. Lee, Phys. Rev. D **78**, 114027 (2008) [arXiv:0805.1989 [hep-ph]].
[29] E. Braaten and J. Lee, Phys. Rev. D **63**, 071501 (2001) [arXiv:hep-ph/0012244].
[30] M. Krämer, Prog. Part. Nucl. Phys. **47**, 141 (2001) [arXiv:hep-ph/0106120].
[31] E. Braaten and T. C. Yuan, Phys. Rev. D **52**, 6627 (1995) [arXiv:hep-ph/9507398].
[32] C. Amsler *et al.* (Particle Data Group), Phys. Lett. B **667**, 1 (2008).